



- aborting outside designated facilities in metropolitan South Africa? *BJOG* 2005; 112: 1236-1242.
3. Morroni C, Moodley J. The role of pregnancy testing in facilitating access to antenatal care and abortion services in South Africa: a cross-sectional study. *BMC Pregnancy and Childbirth* 2006; 6: 26.
 4. Perslev A, Rorbye C, Boesen HC, Norgaard M, Nilas L. Emergency contraception knowledge and use among Danish women requesting termination of pregnancy. *Contraception* 2002; 66: 427-431.
 5. Lee SWH, Wai MFY, Lai LYH, Ho PC. Women's knowledge of and attitudes towards emergency contraception in Hong Kong: questionnaire survey. *Hong Kong Medical Journal* 1999; 5: 349-352.
 6. Smit J, McFadyen L, Beskinska M, et al. Emergency contraception in South Africa: knowledge, attitude and use among public sector primary healthcare clients. *Contraception* 2001; 64: 333-337.

Rape survivors and the provision of HIV post-exposure prophylaxis

S Killian, S Suliman, N Fakier, S Seedat

To the Editor: Rape and HIV/AIDS are two scourges of epidemic proportion in South Africa, which is known for high levels of sexual violence and one of the fastest-growing HIV epidemics in the world.¹ While the link between rape and long-term physical and mental health problems is well established,² the vast majority of rapes go unreported and only a small proportion of women attend health care services after rape, with many believing that their actions will not lead to punishment for the perpetrator.³ Another factor that may worsen matters even further is the latest version of the original bill on sexual offences (Criminal Law (Sexual Offences) Amendment Bill) passed in 2003. The latest version of the Bill states that the provision of post-exposure prophylaxis (PEP) is dependent on the 'victim' laying a charge.⁴ This is likely to have a major negative impact on rape survivors receiving PEP within 72 hours and may impede the ability of medical and police officials to obtain medical evidence that is crucial for the successful prosecution of the perpetrator. Furthermore, the Bill does away with measures to provide for the comprehensive management, care and treatment of rape survivors and the omission of these measures is arguably dismissive of the needs of rape survivors. Another issue of concern is that women who do lay a charge and attend a health care service after rape do not optimally access and utilise the range of services available. Many primary and secondary level services in this country provide HIV testing and counselling, PEP, treatment of sexually

transmitted infections, pregnancy testing and prevention, forensic examination and treatment of injuries. Health services providing the spectrum of post-rape care have an important role to play in changing the horrendous image of rape and HIV/AIDS and more particularly in fulfilling the needs of rape survivors. However, compliance and follow-up remains a major challenge at ground level.

Some of the well-known public health services providing post-rape care in the Western Cape are the Karl Bremer Rape Centre, Groote Schuur Hospital, and the Thuthuzela Rape Centre established at G F Jooste Hospital. We conducted a retrospective analysis of rape survivors presenting to Karl Bremer Rape Centre over a 6-month period (January - June 2005). Three hundred and sixty-three confirmed cases of female rape were identified through chart review, with a mean age of 23.2 years (SD = 10.1). The highest number of patient visits was recorded on Saturdays and Sundays and it may therefore be assumed that the frequency of rape specifically increases over weekends. There was clinical evidence of alcohol use in more than a quarter of rape survivors ($N = 102$) and clinical evidence of drugs in about 2% ($N = 7$). Notably, in 50 reported cases (14%) 2 or more perpetrators were involved, with the number of perpetrators ranging from 1 to 8. The vast majority of perpetrators were not known to the survivor ($N = 112$; 31%), while in 4% of cases ($N = 14$) a familial relationship (the rapist being a parent, sibling, cousin, step-sibling, uncle) existed. More than a third of women had sustained genital ($N = 128$; 35%) or other physical injuries ($N = 93$; 26%). Approximately 13% ($N = 47$) tested HIV positive (on the initial Rapid Test) and 18 patients (5%) tested positive for pregnancy. Perpetrators used a condom in only 6% of cases ($N = 22$). Only 12 women (3.4%) reported for a follow-up assessment at the clinic within a 3-month period.

The findings are significant when considering the potential impact of PEP provision and post-rape care on health services, as well as on police and related services.⁵ The involvement of multiple perpetrators in about a quarter of cases is not

MRC Unit on Anxiety and Stress Disorders, Department of Psychiatry, Stellenbosch University, Tygerberg, W Cape

S Killian, BA Hons (Psychology)

S Suliman, MA (Research Psychology)

S Seedat, MB ChB, MMed (Psych), FCPsych, PhD

Alcohol and Drug Abuse Research Unit, Medical Research Council, Parow, W Cape

N Fakier, MA (Research Psychology)

Corresponding author: S Killian (sanjak@sun.ac.za)



insubstantial when assessing the potential impact of PEP, since rape by multiple perpetrators potentially means multiple HIV-infected perpetrators.⁵ Involvement of multiple perpetrators (particularly in the context of gang rape) is likely to be associated with greater health care needs (physical and emotional) on the part of survivors. We need to be cognisant of this if we are aiming to deliver efficient post-rape care.

Multidisciplinary care services with a holistic approach, such as the Karl Bremer Rape Centre, play a crucial role in addressing the various needs of rape survivors. A recent South African study on women's experiences of, and preferences for, services after rape clearly indicated that rape survivors have a need for holistic post-rape care.⁶ The study showed that women particularly valued the availability of HIV prophylaxis and a sensitive health care provider who could provide counselling, and that women were willing to travel further to clinics for counselling, rigorous examination and HIV prophylaxis. That said, the long-term success of health services in preventing HIV/AIDS and in fulfilling the specific needs of rape survivors may be undermined by a lack of research on the effectiveness of PEP following sexual assault. Research on the latter needs

to be done together with ongoing efforts to operationalise and evaluate what is currently available and applicable within the South African context.⁵ Empirical data on the efficacy of PEP will essentially contribute to the provision of more optimal services for rape survivors. For many rape survivors in South Africa, rape means a death sentence. More efficient services as well as increased conviction rates for perpetrators are concrete ways of addressing criminal sexual assault and will ultimately go a long way toward preventing new HIV infections.⁷

1. Kim JC, Martin LJ, Denny L. Rape and HIV post-exposure prophylaxis: addressing the dual epidemics in South Africa. *Reprod Health Matters* 2003; 11(22): 101-112.
2. Campbell R, Wasco SM. Understanding rape and sexual assault: 20 years of progress and future directions. *J Interpers Violence* 2005; 20(1): 127-131.
3. Jewkes R, Abrahams N. The epidemiology of rape and sexual coercion in South Africa: an overview. *Soc Sci Med* 2002; 55(7): 1231-1244.
4. Criminal Law (Sexual Offences) Amendment Bill, 2006, Republic of South Africa.
5. Kim JC. Rape and HIV post-exposure prophylaxis: its relevance and the reality in South Africa. In: *Violence Against Women and HIV/AIDS: Setting the Research Agenda*. Geneva: World Health Organization, 2001: 38-41. WHO/FCH/GWH/01.08.
6. Christofides NJ, Muirhead D, Jewkes RK, Penn-Kekana L, Conco DN. Women's experiences of and preferences for service after rape in South Africa: interview study. *BMJ* 2006; 332: 209-213.
7. Kalichman SC, Simbayi LC. Sexual assault history and risks for sexually transmitted infections among women in an African township in Cape Town, South Africa. *AIDS Care* 2007; 16(6): 681-689.

amcsa 2007
Aesthetic Medicine Congress South Africa

Aesthetic Medicine Congress South Africa
5 - 6 October 2007 The Estate Centre, Gallagher Estate, Gauteng

The second Aesthetic Medicine Congress of South Africa will be held at The Estate Centre, Gallagher Estate. The purpose of the congress is to disseminate to a South African Medical professional audience the latest teachings on aesthetic and anti-ageing treatment modalities. The primary focus being on global guiding principles for the treatment of your patient. The congress will cover a range of topics in the field of Aesthetic Medicine and feature comprehensive and leading edge information through a dynamic scientific program and high quality speakers.

The scientific program will cover the following important topics:

- **Botox and Fillers:** 2007 Update, advance techniques and live workshops with top international speakers;
- **Peeling:** From superficial to deep peeling indications and complications;
- **Mesotherapy:** Scientific approaches, research and literature reviews;
- **Lasers and Lights:** Recently launched technology;
- **Anti-ageing Medicine:** How to postpone skin as well as body ageing;
- **NEW session on sexual ageing;**
- **Sclerotherapy:** Workshops and presentations;
- **Aesthetic practice management and marketing;**
- **Non-surgical Lipolysis:** Latest updates and devices;
- **Medico-legal aspects in South Africa;**
- **Skin Rejuvenation, Skin ageing, Cosmetology and Acne Management.**

International and local professionals, who are experts in their fields of Aesthetic Medicine, will present up to date information. They include: Dr Hervé Raspaldo, Dr Cedric Kron, Debra Robson Lawrence, Prof Max Lafontan, Dr Sohaïl Mansoor, Dr Mike Perring, Oded Rose, Dr Anders Strand, Darren Thomas, Dr Marlene Wasserman, Prof Ronald W Pero, Mr Chris J Ingfield (BSc, MBBS, FRCS(Plast)), Dr Ilse du Plessis, Ingrid du Plessis, Brink Eksteen, Dr Des Fernandes, Dr Nardus le Grange, Dr Geraldine Mitton, Brent Murphy, Dr Annemarie Potgieter, Dr Riekje Smit, Prof J.R. Snyman, Dr Bradley Wagemaker, Dr Vivien Jandera.

The Congress will also include an exhibition featuring the latest in cutting edge products and technology.

In association with:



For more information contact us at 012 543 9547 or visit www.aestheticcongress.co.za